

seminarium Matematyka Dyskretna

wtorek, 6 grudnia 2022 r., godz. 12:30, s. 304 A3-A4

Maximal partial designs

Mariusz Meszka WMS AGH

A few problems related to various combinatorial designs in terms of their maximality will be discussed.

One of the most examined cases concerns maximal partial latin squares. A *partial* latin square of order n is an $n \times n$ array in which each cell is either empty or contains a single symbol from an n-element set S such that each symbol occurs at most once in each row and at most once in each column. A partial latin square is *maximal* if no empty cell can be filled with an element of S without violating latin conditions.

Evidently, a partial latin square of order n corresponds to a proper edge-coloring of a balanced bipartite graph H = (V, U, E) with at most $n = \chi'(K_{n,n})$ colors, where |U| = |V| = n. In this way considering maximal edge-colorings of non-bipartite graphs makes an obvious next step. Let G be a graph of order n. A maximal edge-coloring of G is a proper edge-coloring with $\chi'(K_n)$ colors such that no edge of the complement \overline{G} can be attached to G without violating conditions of proper edge-coloring. For given n, a spectrum MEC(n) is defined to be the set of all sizes of graphs of order n which admit maximal edge-colorings.

Another class of objects, strongly related to edge-colorings of graphs, are partial Room squares. A *partial Room square* of order n and side n - 1 on an n-element set S is an $(n-1) \times (n-1)$ array F satisfying the following properties:

(1) every cell of F is either empty or contains an unordered pair of symbols from S,

(2) every symbol of S occurs at most once in each row and at most once in each column of F,

(3) every unordered pair of symbols of S occurs in at most one cell of F.

A partial Room square is *maximal* if no further pair of elements can be placed into any unoccupied cell without violating the conditions that define a partial Room square. A *spectrum* MPRS(n) is the set of volumes of maximal partial Room squares of order n, where the *volume* means the number of occupied cells.

In all of these cases, the common aim is to determine spectra.