

seminarium Matematyka Dyskretna

wtorek, 5 marca 2024 r., godz. 12:30, s. 612 C
7 $\,$

3-colourability, diamonds and spindle graphs

Ingo Schiermeyer TU Freiberg (joint work with Christoph Brause and Přemysl Holub)

The 3-colourability problem is an NP-complete problem which remains NP-complete for claw-free graphs and for graphs with maximum degree four. In this talk we will consider induced subgraphs, among them are the *claw* $(K_{1,3})$, the *bull* (a triangle with two pendent edges), the *net* (a triangle with three pendent edges), and the *diamond* (the graph $K_4 - e$).

Our main result is a complete characterization of all 3-colourable (claw, H)-free graphs for $H \in \{bull, net, P_7\}$. We will present a description of all non 3-colourable (claw, H)-free graphs for $H \in \{bull, net, P_7\}$ in terms of diamonds. Moreover, we will show extensions of this characterization to larger graph classes by taking supergraphs of the claw or the bull.

- [1] B. Randerath and I. Schiermeyer, Vertex colouring and forbidden subgraphs a survey, Graphs and Combinatorics 20 (2004) 1-40.
- [2] I. Schiermeyer and B. Randerath, Polynomial χ-Binding Functions and Forbidden Induced Subgraphs: A Survey, Graphs and Combinatorics 35 (1) (2019) 1–31.