



SEMINARIUM MATEMATYKA DYSKRETNA

wtorek, 25 marca 2025 r., godz. 12:30, s. 612 C7

Weak and strong local irregularity of digraphs

Igor Grzelec
WMS AGH

Local Irregularity Conjecture proposed by Baudon, Bensmail, Przybyło and Woźniak in 2015 states that every connected graph, except special cacti, can be decomposed into at most three *locally irregular graphs*, i.e., graphs in which adjacent vertices have different degrees. The concept of local irregularity was defined for digraphs in several different ways. After short introduction about the above mentioned Local Irregularity Conjecture we present the already known concepts of local irregularity for digraphs with motivations, main conjectures and known results. Then we introduce the following new methods of defining a *locally irregular digraph*. The first one, *weak local irregularity*, is based on distinguishing adjacent vertices by *indegree-outdegree pairs*, and the second one, *strong local irregularity*, asks for *different balanced degrees* (i.e., difference between the outdegree and the indegree of a vertex) of adjacent vertices. For both of these irregularities, we define locally irregular decompositions and colorings of digraphs. We also provide related conjectures on the minimum number of colors in weak and strong locally irregular colorings and support them with new results for various classes of digraphs.

This is a joint work Alfréd Onderko and Mariusz Woźniak.